

U.S. Preventive Services Task Force

Grade Definitions

The U.S. Preventive Services Task Force (USPSTF) assigns one of five letter grades (A, B, C, D, or I). The USPSTF changed its grade definitions based on a change in methods in May 2007 and again in July 2012, when it updated the definition of and suggestions for practice for the grade C recommendation.

Grade Definitions After July 2012

What the Grades Mean and Suggestions for Practice

The USPSTF updated its definition of and suggestions for practice for the grade C recommendation. This new definition applies to USPSTF recommendations voted on after July 2012. Describing the strength of a recommendation is an important part of communicating its importance to clinicians and other users. Although most of the grade definitions have evolved since the USPSTF first began, none has changed more noticeably than the definition of a C recommendation, which has undergone three major revisions since 1998. Despite these revisions, the essence of the C recommendation has remained consistent: at the population level, the balance of benefits and harms is very close, and the magnitude of net benefit is small. Given this small net benefit, the USPSTF has either not made a recommendation “for or against routinely” providing the service (1998), recommended “against routinely” providing the service (2007), or recommended “selectively” providing the service (2012). Grade C recommendations are particularly sensitive to patient values and circumstances. Determining whether or not the service should be offered or provided to an individual patient will typically require an informed conversation between the clinician and patient.

Grade	Definition	Suggestions for Practice
A	The USPSTF recommends the service. There is high certainty that the net benefit is substantial.	Offer or provide this service.
B	The USPSTF recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial.	Offer or provide this service.
C	The USPSTF recommends selectively offering or providing this service to individual patients based on professional judgment and patient preferences. There is at least moderate certainty that the net benefit is small.	Offer or provide this service for selected patients depending on individual circumstances.
D	The USPSTF recommends against the service. There is moderate or high certainty that the service has no net benefit or that the harms outweigh the benefits.	Discourage the use of this service.
I Statement	The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined.	Read the clinical considerations section of USPSTF Recommendation Statement. If the service is offered, patients should understand the uncertainty about the balance of benefits and harms.

Levels of Certainty Regarding Net Benefit

Level of Certainty*	Description
High	The available evidence usually includes consistent results from well-designed, well-conducted studies in representative primary care populations. These studies assess the effects of the preventive service on health outcomes. This conclusion is therefore unlikely to be strongly affected by the results of future studies.
Moderate	The available evidence is sufficient to determine the effects of the preventive service on health outcomes, but confidence in the estimate is constrained by such factors as: <ul style="list-style-type: none"> • The number, size, or quality of individual studies. • Inconsistency of findings across individual studies. • Limited generalizability of findings to routine primary care practice. • Lack of coherence in the chain of evidence. As more information becomes available, the magnitude or direction of the observed effect could change, and this change may be large enough to alter the conclusion.
Low	The available evidence is insufficient to assess effects on health outcomes. Evidence is insufficient because of: <ul style="list-style-type: none"> • The limited number or size of studies. • Important flaws in study design or methods. • Inconsistency of findings across individual studies. • Gaps in the chain of evidence. • Findings not generalizable to routine primary care practice. • Lack of information on important health outcomes. More information may allow estimation of effects on health outcomes.

* The USPSTF defines certainty as "likelihood that the USPSTF assessment of the net benefit of a preventive service is correct." The net benefit is defined as benefit minus harm of the preventive service as implemented in a general, primary care population. The USPSTF assigns a certainty level based on the nature of the overall evidence available to assess the net benefit of a preventive service.

Retrieved from the USPSTF Web Site on May 13, 2018
<https://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions>

#